

Application Ser. No.: 10/613,588


Filing Date: July 3, 2003

Examiner: Huang, Evelyn Mei

Amendment Pursuant to 37 C.F.R. § 1.121IN THE CLAIMS:

The claims set forth below with amendments as indicated will replace all prior versions and listing of claims in the application.

1. (currently amended) A compound of the formula I



(I)

or a stereoisomeric form or a pharmaceutically acceptable salt of the compound of the formula I, wherein

A is $-(C_1-C_6)$ -alkyl, in which alkyl is straight-chain or branched and is optionally substituted, once or more, independently of each other, by

$-O-R^1$ or

$-C(O)-OR^1$,

$-C(O)-NR^1R^1$,

$-C(O)-NR^1-SO_2R^1$,

$-NR^1R^1$,

$-CN$, in which R^1 is

hydrogen,

$-(C_1-C_6)$ -alkyl,

$-(C_6-C_{14})$ aryl or

fluoroalkyl of the formula $-C_nH_xF_y$ or fluoroalkoxy of the formula $-OC_nH_xF_y$, wherein n is an integer from 1 to 6, x is an integer from 0 to 12, y is an integer from 1 to 13 and sum of x and y is $2n + 1$,

$-O-R^1$,

$-SR^1$,

Application Ser. No.: 10/613,588

Filing Date: July 3, 2003

Examiner: Huang, Evelyn Mei

-S(O)-R¹

-S(O)₂-R¹

-C(O)-OR¹,

fluoroalkyl of the formula -C_nH_xF_y or fluoroalkoxy of the formula

-OC_nH_xF_y, wherein n is an integer from 1 to 6, x is an integer from 0 to

12, y is an integer from 1 to 13 and sum of x and y is 2n + 1,

-C(O)-NR¹R¹,

-C(O)-NR¹-SO₂R¹,

-NR¹R¹,

-CN,

monocyclic or polycyclic heteroaryl having from 5 to 14 ring members, which contains 1 to 5 heteroatoms as ring members, wherein heteroatoms are selected from N, O and S, and in which heteroaryl is unsubstituted or optionally substituted once or more, independently of each other, by R², in which R² is

-(C₁-C₄)-alkyl,

-OH,

-O-(C₁-C₄)-alkyl,

halogen,

-N(R³)-R⁴, in which R³ and R⁴ are, independently of each other, hydrogen atom or -(C₁-C₄)-alkyl,

fluoroalkyl of the formula -C_nH_xF_y or fluoroalkoxy of the formula -OC_nH_xF_y, wherein n is an integer from 1 to 4, x is an integer from 0 to 8, y is an integer from 1 to 9 and sum of x and y is 2n + 1,

-CN,

-SR¹,

-S(O)-R¹,

-S(O)₂-R¹ or

-C(O)-NR¹R¹,

-(C₃-C₆)-cycloalkyl, in which cycloalkyl is unsubstituted or substituted, once or more, independently of each other, by R², and R² is defined as above,

monocyclic or bicyclic heterocycle having from 5 to 12 ring members, which contains 1 to 5 heteroatoms as ring members, wherein

Application Ser. No.: 10/613,588

Filing Date: July 3, 2003

Examiner: Huang, Evelyn Mei

heteroatoms are selected from N, O and S, and in which heterocycle is unsubstituted or substituted, once or more, independently of each other, by R², and R² is defined as above,

B is a covalent bond,

-C=CR¹-,

—C≡C—

-O(CH₂)_a-, in which a is an integer from 1 to 4,
O, S, NR²-, -C(O)-, -NR²-C(O)-, -C(O)-NR²-, -NR²-SO₂-,
-SO₂-NR²-, -NR²-C(O)-NR²-, and R² is defined as above, or
-(C₁-C₄)-alkylene, in which alkylene is straight-chain or
branched and is optionally substituted, once or more,
independently of each other, by R¹, and R¹ is defined as
above,

D is -(C₁-C₆)-alkyl, in which alkyl is straight-chain or branched
and is optionally substituted, once or more, independently of each
other, by R¹, and R¹ is defined as above,

monocyclic or polycyclic heteroaryl having from 5 to 14 ring
members, which contains 1 to 5 heteroatoms as ring
members, and in which heteroaryl is unsubstituted or is
substituted once or more, independently of each other, by
R², and R² is defined as above,

monocyclic or bicyclic heterocycle having from 5 to 12 ring
members, which contains 1 to 5 heteroatoms as ring
members, and in which heterocycle is unsubstituted or
substituted, once or more, independently of each other, by
R², and R² is defined as above,

-(C₆-C₁₄)-aryl, in which aryl is unsubstituted or substituted,
once or more, independently of each other, by R², and R² is
defined as above, or

-(C₃-C₆)-cycloalkyl, in which cycloalkyl is unsubstituted or
substituted, once or more, independently of each other, by
R², and R² is defined as above, or

B-D is hydrogen,
halogen,

fluoroalkyl of the formula -C_nH_xF_y or fluoroalkoxy of the
formula -OC_nH_xF_y, wherein n is an integer from 1 to 4, x is

Application Ser. No.: 10/613,588

Filing Date: July 3, 2003

Examiner: Huang, Evelyn Mei

an integer from 0 to 8, y is an integer from 1 to 9 and sum of x and y is $2n + 1$,

$-(CH_2)_a-Y-R^3$, in which a is an integer from 1 to 4, Y is O, S, NR^2 , and R^3 is

$-(C_1-C_6)$ -alkyl,
 $-(C_6-C_{14})$ -aryl,
 $-(C_3-C_6)$ -cycloalkyl, and

R is hydrogen,
 $-(C_1-C_6)$ -alkyl, or
 $-(C_6-C_{14})$ -aryl- $-(C_1-C_6)$ -alkyl, in which aryl is unsubstituted or substituted, once or more, independently of each other, by R^2 , and R^2 is defined as above, and

X and Z are identical or different and are, independently of each other selected from:

hydrogen atom,
 $-(C_1-C_4)$ -alkyl,
 $-OH$,
 $-O-(C_1-C_4)$ -alkyl),
halogen,
fluoroalkyl of the formula $-C_nH_xF_y$ or fluoroalkoxy of the formula $-OC_nH_xF_y$, wherein n is an integer from 1 to 6, x is an integer from 0 to 12, y is an integer from 1 to 13 and sum of x and y is $2n + 1$,
 $-C(O)-OR^1$,
 $-C(O)-NR^1R^1$,
 $-C(O)-NR^1-SO_2R^1$,
 $-NR^1R^1$,
 $-NR^1-C(O)-NR^1R^1$,
 $-NR^1-C(O)-R^1$,
 $-NR^1-C(O)-OR^1$,
 $-O-C(O)-NR^1R^1$,
 $-CN$,
 $-SR^1$,
 $-S(O)-R^1$,
 $-S(O)_2-R^1$,
 $-S(O)_2-NR^1R^1$,
 $-NR^1-SO_2-R^1$, in which R^1 is as defined above,

Application Ser. No.: 10/613,588

Filing Date: July 3, 2003

Examiner: Huang, Evelyn Mei

monocyclic or bicyclic heterocycle having from 5 to 12 ring members, which contains 1 to 5 heteroatoms as ring members, wherein heteroatoms are selected from N, O and S, and in which heterocycle is unsubstituted or substituted, once or more, independently of each other, by R², and R² is defined as above, or

-(C₃-C₆)-cycloalkyl, in which cycloalkyl is unsubstituted or substituted, once or more, independently of each other, by R², and R² is defined as above,

with the proviso that when A is -(C₁-C₆)-alkyl, in which alkyl is straight-chain or branched and is optionally substituted, once or more, independently of each other, by

-O-R¹ or

-C(O)-OR¹, in which R¹ is

hydrogen,

-(C₁-C₆)-alkyl,

-O-R¹, in which R¹ is as defined in this proviso,

-C(O)-OR¹, in which R¹ is as defined in this proviso, or

monocyclic or polycyclic heteroaryl having from 5 to 14 ring members, which contains 1 to 5 heteroatoms as ring members, wherein heteroatoms are selected from N, O and S, and in which heteroaryl is unsubstituted or optionally substituted once or more, independently of each other, by R², in which R² is

hydrogen,

-(C₁-C₄)-alkyl,

-OH,

-O-(C₁-C₄)-alkyl,

halogen, or

-N(R³)-R⁴, in which R³ and R⁴ are, independently of each other, hydrogen atom or -(C₁-C₄)-alkyl, then at least one of the following applies:

B is not a covalent bond or -(C₁-C₄)-alkylene,

D is not heteroaryl, heterocycle, -(C₆-C₁₄)-aryl, (C₃-C₆)-cycloalkyl, or

X and Z are not -(C₁-C₄)-alkyl, -OH, -O-(C₁-C₄)-alkyl, or halogen.

2. (original) A compound of the formula I as claimed in claim 1, wherein

A is -(C₁-C₃)-alkyl, in which alkyl is straight-chain or branched and is optionally substituted, once or more, independently of each other, by

Application Ser. No.: 10/613,588

Filing Date: July 3, 2003

Examiner: Huang, Evelyn Mei

-O-R¹, or
 -C(O)-OR¹, in which R¹ is
 hydrogen,
 -(C₁-C₃)-alkyl, or
 -CF₃

fluoroalkyl of the formula -C_nH_xF_y or fluoroalkoxy of the formula
 -OC_nH_xF_y, wherein n is an integer from 1 to 3, x is an integer from 0
 to 6, y is an integer from 1 to 7 and sum of x and y is 2n + 1,

B is a covalent bond or O,

D is phenyl or naphthyl, in which phenyl or naphthyl is unsubstituted
 or substituted, once or more, independently of each other, by R², in
 which R² is

fluorine, chlorine or bromine,
 -OH,
 -CF₃,
 -SR¹, in which R¹ is defined as above,
 -(C₁-C₄)-alkyl
 -O-(C₁-C₂)-alkyl or
 -N(R³)-R⁴, in which R³ and R⁴ are, independently of

each other, hydrogen atom or -(C₁-C₃)-alkyl,

heteroaryl selected from the group consisting of
 pyridyl, furanyl, pyrrolyl, isoxazolyl, benzofuranyl, benzothiophenyl,
 quinolinyl, isoquinolinyl, quinoxaliny and thiophenyl, in which
 heteroaryl is unsubstituted or substituted, once or more,
 independently of each other, by R², in which R² is defined as above
 or

-(C₄-C₆)-cycloalkyl, in which cycloalkyl is
 unsubstituted or substituted, once or more, independently of each
 other, by R², and R² is defined as above, or

B-D is ((CH₂)_a-Y-R³, in which a is an integer from 1 to 2, Y is O and R³
 is -(C₁-C₃)-alkyl, and

R is hydrogen,

-(C₁-C₃)-alkyl, or
 -phenyl-(C₁-C₃)-alkyl, and

X and Z are identical or different and are, independently of each
 other, hydrogen, -C(O)-O(C₁-C₃)-alkyl, -OCH₃, -N(CH₃)₂ or halogen.

3. (original) A compound of the formula 1 as claimed in claim 1, wherein the
 compound of the formula I is selected from the group consisting of:

Application Ser. No.: 10/613,588

Filing Date: July 3, 2003

Examiner: Huang, Evelyn Mei

5-pyridin-2-yl-3-trifluoromethyl-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(2-trifluoromethyl-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(3-trifluoromethyl-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(4-trifluoromethyl-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
1,3-dimethyl-5-(3-trifluoromethylphenyl)-1H-pyrazolo[4,3-c]-isoquinoline,
5-phenyl-3-trifluoromethyl-1H-pyrazolo[4,3-c]isoquinoline,
1,3-dimethyl-5-(3-trifluoromethylphenyl)-1H-pyrazolo[4,3-c]-isoquinoline,
1,3-dimethyl-5-(2,6-difluorophenyl)-1H-pyrazolo[4,3-c]-isoquinoline,
1-benzyl-5-cyclohexyl-3-methyl-1H-pyrazolo[4,3-c]-isoquinoline,
1-benzyl-5-naphthyl-3-methyl-1H-pyrazolo[4,3-c]-isoquinoline,
5-methoxymethyl-3-methyl-1H-pyrazolo[4,3-c]-isoquinoline,
7-methoxycarbonyl-3-methyl-5-phenyl-1H-pyrazolo[4,3-c]-isoquinoline,
7-methoxycarbonyl-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]-
isoquinoline,
7-dimethylamino-3-methyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,
7-dimethylamino-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]-isoquinoline,
6-dimethylamino-3-methyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,
6-dimethylamino-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]-isoquinoline,
8-dimethylamino-3-methyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,
8-dimethylamino-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]-isoquinoline,
1,3-dimethyl-5-(3-methyl-thiophen-2-yl)-1H-pyrazolo[4,3-c]-isoquinoline,
3-methyl-5-phenyl-9-trifluoromethyl-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-pyridin-2-yl-9-trifluoromethyl-1H-pyrazolo[4,3-c]-isoquinoline,
and
3-methyl-5-(2,3,4,5,6-pentafluoro-phenyl)-1H-pyrazolo[4,3-c]-isoquinoline.

4. (currently amended) A compound selected from the group consisting of:

5-benzo[b]thiophene-2yl-3-methyl-1H-pyrazolo[4,3-c]isoquinoline,
7-bromo-3-methyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,
7-bromo-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]isoquinoline,

Application Ser. No.: 10/613,588

Filing Date: July 3, 2003

Examiner: Huang, Evelyn Mei

5-(3-chloro-benzo[b]thiophen-2-yl)-3-methyl-1H-pyrazolo[4,3-c]-isoquinoline,
5-(2-chloro-pyridin-3-yl)-3-methyl-1H-pyrazolo[4,3-c]isoquinoline,
5-(6-chloro-pyridin-3-yl)-3-methyl-1H-pyrazolo[4,3-c]isoquinoline,
3-ethyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,
3-ethyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]isoquinoline,
3-ethyl-5-pyridin-3-yl-1H-pyrazolo[4,3-c]isoquinoline,
5-furan-2-yl-3-methyl-1H-pyrazolo[4,3-c]isoquinoline,
7-methoxy-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(3-methyl-benzofuran-2-yl)-1H-pyrazolo[4,3-c]-isoquinoline,
~~3-methyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,~~
6-chloro-3-methyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,
6-fluoro-3-methyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,
8-chloro-3-methyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,
8-fluoro-3-methyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,
9-chloro-3-methyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,
9-fluoro-3-methyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,
3,9-dimethyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(2-methyl-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(3-methyl-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
~~3-methyl-5-(4-methyl-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,~~
3-methyl-5-(2-bromo-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(3-bromo-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(2-chloro-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
~~3-methyl-5-(4-chloro-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,~~
3-methyl-5-(2,4-dichloro-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(3,4-dichloro-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(2,6-dichloro-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(2-fluoro-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(4-fluoro-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,

Application Ser. No.: 10/613,588

Filing Date: July 3, 2003

Examiner: Huang, Evelyn Mei

3-methyl-5-(2,4-difluoro-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(2,6-difluoro-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
6-chloro-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]isoquinoline,
6-fluoro-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]isoquinoline,
8-chloro-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]isoquinoline,
8-fluoro-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]isoquinoline,
9-chloro-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]isoquinoline,
9-fluoro-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]isoquinoline,
3,9-dimethyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(1-methyl-1H-pyrrol-2-yl)-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-quinolin-2-yl-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-quinolin-3-yl-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-quinoxalin-2-yl-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-thiophen-2-yl-1H-pyrazolo[4,3-c]isoquinoline, and
3-methyl-5-(3-methyl-thiophen-2-yl)-1H-pyrazolo[4,3-c]isoquinoline.

5. (original) A pharmaceutical composition comprising a therapeutically effective content of at least one compound of the formula I as claimed in claim 1 together with a pharmaceutically suitable carrier optionally in combination with a suitable additive, other active compounds and auxiliary substances.
6. (original) A method of treating a disease condition associated with the increased activity of NIK comprising administering to a patient suffering from said disease condition a therapeutically effective amount of a compound according to claim 1.
7. (original) The method as claimed in claim 6, wherein the compound is according to claim 2.

Application Ser. No.: 10/613,588

Filing Date: July 3, 2003

Examiner: Huang, Evelyn Mei

8. (original) The method as claimed in claim 6 wherein said compound is selected from the group consisting of:
- 5-pyridin-2-yl-3-trifluoromethyl-1H-pyrazolo[4,3-c]isoquinoline,
 - 3-methyl-5-(2-trifluoromethyl-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
 - 3-methyl-5-(3-trifluoromethyl-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
 - 3-methyl-5-(4-trifluoromethyl-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
 - 1,3-dimethyl-5-(3-trifluoromethylphenyl)-1H-pyrazolo[4,3-c]-isoquinoline,
 - 5-phenyl-3-trifluoromethyl-1H-pyrazolo[4,3-c]isoquinoline,
 - 1,3-dimethyl-5-(3-trifluoromethylphenyl)-1H-pyrazolo[4,3-c]-isoquinoline,
 - 1,3-dimethyl-5-(2,6-difluorophenyl)-1H-pyrazolo[4,3-c]-isoquinoline,
 - 1-benzyl-5-cyclohexyl-3-methyl-1H-pyrazolo[4,3-c]-isoquinoline,
 - 1-benzyl-5-naphthyl-3-methyl-1H-pyrazolo[4,3-c]-isoquinoline,
 - 5-methoxymethyl-3-methyl-1H-pyrazolo[4,3-c]-isoquinoline,
 - 7-methoxycarbonyl-3-methyl-5-phenyl-1H-pyrazolo[4,3-c]-isoquinoline,
 - 7-methoxycarbonyl-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]-isoquinoline,
 - 7-dimethylamino-3-methyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,
 - 7-dimethylamino-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]-isoquinoline,
 - 1,3-dimethyl-5-(3-methyl-thiophen-2-yl)-1H-pyrazolo[4,3-c]-isoquinoline,
 - 3-methyl-5-phenyl-9-trifluoromethyl-1H-pyrazolo[4,3-c]isoquinoline,
 - 3-methyl-5-pyridin-2-yl-9-trifluoromethyl-1H-pyrazolo[4,3-c]-isoquinoline,
 - and
 - 3-methyl-5-(2,3,4,5,6-pentafluoro-phenyl)-1H-pyrazolo[4,3-c]-isoquinoline.
9. (original) The method as claimed in claim 6, wherein the disease condition is caused due to an inflammatory component.
10. (original) The method as claimed in claim 6, wherein the diseases are osteoarthritis, rheumatoid arthritis, asthma, irritable bowel disease, Alzheimer's disease, stroke, diabetes, atherosclerosis, multiple sclerosis,


Application Ser. No.: 10/613,588

Filing Date: July 3, 2003

Examiner: Huang, Evelyn Mei

rejection reactions on the part of the body against a transplanted organ or
rejection reactions on the part of the transplanted organ against the body.

11. (currently amended) A pharmaceutical composition comprising a compound
of the formula (I)



(I)

or a stereoisomeric form or a pharmaceutically acceptable salt of the
compound of the formula I, wherein

A is $-(C_1-C_6)\text{-alkyl}$, in which alkyl is straight-chain or branched
and is optionally substituted, once or more, independently of each
other, by

$-O-R^1$ or

$-C(O)-OR^1$,

$-C(O)-NR^1R^1$,

$-C(O)-NR^1-SO_2R^1$,

$-NR^1R^1$,

$-CN$, in which R^1 is

hydrogen,

$-(C_1-C_6)\text{-alkyl}$,

$-(C_6-C_{14})\text{aryl}$ or

fluoroalkyl of the formula $-C_nH_xF_y$ or fluoroalkoxy of

the formula $-OC_nH_xF_y$, wherein n is an integer from 1

Application Ser. No.: 10/613,588

Filing Date: July 3, 2003

Examiner: Huang, Evelyn Mei

to 6, x is an integer from 0 to 12, y is an integer from 1 to 13 and sum of x and y is $2n + 1$,

-O-R¹,

-SR¹,

-S(O)-R¹

-S(O)₂-R¹

-C(O)-OR¹,

fluoroalkyl of the formula -C_nH_xF_y or fluoroalkoxy of the formula -OC_nH_xF_y, wherein n is an integer from 1 to 6, x is an integer from 0 to 12, y is an integer from 1 to 13 and sum of x and y is $2n + 1$,

-C(O)-NR¹R¹,

-C(O)-NR¹-SO₂R¹,

-NR¹R¹,

-CN,

monocyclic or polycyclic heteroaryl having from 5 to 14 ring members, which contains 1 to 5 heteroatoms as ring members, wherein heteroatoms are selected from N, O and S, and in which heteroaryl is unsubstituted or optionally substituted once or more, independently of each other, by R², in which R² is

-(C₁-C₄)-alkyl,

-OH,

-O-(C₁-C₄)-alkyl,

halogen,

-N(R³)-R⁴, in which R³ and R⁴ are, independently of each other, hydrogen atom or -(C₁-C₄)-alkyl,

fluoroalkyl of the formula -C_nH_xF_y or fluoroalkoxy of the formula -OC_nH_xF_y, wherein n is an integer from 1 to 4, x is an

Application Ser. No.: 10/613,588

Filing Date: July 3, 2003

Examiner: Huang, Evelyn Mei

integer from 0 to 8, y is an integer from 1 to 9 and sum of x and y is $2n + 1$,

-CN,

-SR¹,

-S(O)-R¹,

-S(O)₂-R¹ or

-C(O)-NR¹R¹,

-(C₃-C₈)-cycloalkyl, in which cycloalkyl is unsubstituted or substituted, once or more, independently of each other, by R², and R² is defined as above,

monocyclic or bicyclic heterocycle having from 5 to 12 ring members, which contains 1 to 5 heteroatoms as ring members, wherein heteroatoms are selected from N, O and S, and in which heterocycle is unsubstituted or substituted, once or more, independently of each other, by R², and R² is defined as above,

B is a covalent bond,

-C=CR¹-,

—C≡C— ,

-O(CH₂)_a-, in which a is an integer from 1 to 4,

O, S, NR², -C(O)-, -NR²-C(O)-, -C(O)-NR²-, -NR²-SO₂-, -SO₂-NR²-, -NR²-C(O)-NR²-, and R² is defined as above, or

-(C₁-C₄)-alkylene, in which alkylene is straight-chain or branched and is optionally substituted, once or more, independently of each other, by R¹, and R¹ is defined as above,

Application Ser. No.: 10/613,588

Filing Date: July 3, 2003

Examiner: Huang, Evelyn Mei

D is $-(C_1-C_8)$ -alkyl, in which alkyl is straight-chain or branched and is optionally substituted, once or more, independently of each other, by R^1 , and R^1 is defined as above,

monocyclic or polycyclic heteroaryl having from 5 to 14 ring members, which contains 1 to 5 heteroatoms as ring members, wherein heteroatoms are selected from N, O and S, and in which heteroaryl is unsubstituted or is substituted once or more, independently of each other, by R^2 , and R^2 is defined as above,

monocyclic or bicyclic heterocycle having from 5 to 12 ring members, which contains 1 to 5 heteroatoms as ring members, wherein heteroatoms are selected from N, O and S, and in which heterocycle is unsubstituted or substituted, once or more, independently of each other, by R^2 , and R^2 is defined as above,

$-(C_6-C_{14})$ -aryl, in which aryl is unsubstituted or substituted, once or more, independently of each other, by R^2 , and R^2 is defined as above, or

$-(C_3-C_6)$ -cycloalkyl, in which cycloalkyl is unsubstituted or substituted, once or more, independently of each other, by R^2 , and R^2 is defined as above, or

B-D is hydrogen,
halogen,

fluoroalkyl of the formula $-C_nH_xF_y$ or fluoroalkoxy of the formula $-OC_nH_xF_y$, wherein n is an integer from 1 to 4, x is an integer from 0 to 8, y is an integer from 1 to 9 and sum of x and y is $2n + 1$,

Application Ser. No.: 10/613,588

Filing Date: July 3, 2003

Examiner: Huang, Evelyn Mei

$-(CH_2)_a-Y-R^3$, in which a is an integer from 1 to 4, Y is O, S, NR^2 , and R^3 is

$-(C_1-C_6)$ -alkyl,

$-(C_6-C_{14})$ -aryl,

$-(C_3-C_6)$ -cycloalkyl, and

R is hydrogen,
 $-(C_1-C_6)$ -alkyl, or
 $-(C_6-C_{14})$ -aryl- $-(C_1-C_6)$ -alkyl, in which aryl is unsubstituted or substituted, once or more, independently of each other, by R^2 , and R^2 is defined as above, and

X and Z are identical or different and are, independently of each other selected from:

hydrogen atom,

$-(C_1-C_4)$ -alkyl,

$-OH$,

$-O-(C_1-C_4)$ -alkyl),

halogen,

fluoroalkyl of the formula $-C_nH_xF_y$ or fluoroalkoxy of the formula $-OC_nH_xF_y$, wherein n is an integer from 1 to 6, x is an integer from 0 to 12, y is an integer from 1 to 13 and sum of x and y is $2n + 1$.

$-C(O)-OR^1$,

$-C(O)-NR^1R^1$,

$-C(O)-NR^1-SO_2R^1$,

$-NR^1R^1$,

$-NR^1-C(O)-NR^1R^1$,

$-NR^1-C(O)-R^1$,

$-NR^1-C(O)-OR^1$,

$-O-C(O)-NR^1R^1$.

Application Ser. No.: 10/613,588

Filing Date: July 3, 2003

Examiner: Huang, Evelyn Mei

-CN,

-SR¹,

-S(O)-R¹,

-S(O)₂-R¹,

-S(O)₂-NR¹R¹,

-NR¹-SO₂-R¹, in which R¹ is as defined above,

monocyclic or bicyclic heterocycle having from 5 to 12 ring members, which contains 1 to 5 heteroatoms as ring members, wherein heteroatoms are selected from N, O and S, and in which heterocycle is unsubstituted or substituted, once or more, independently of each other, by R², and R² is defined as above, or

-(C₃-C₆)-cycloalkyl, in which cycloalkyl is unsubstituted or substituted, once or more, independently of each other, by R², and R² is defined as above,

with the proviso that when A is -(C₁-C₆)-alkyl, in which alkyl is straight-chain or branched and is optionally substituted, once or more, independently of each other, by

-O-R¹ or

-C(O)-OR¹, in which R¹ is
hydrogen,

-(C₁-C₆)-alkyl,

-O-R¹, in which R¹ is as defined in this proviso,

-C(O)-OR¹, in which R¹ is as defined in this proviso, or

monocyclic or polycyclic heteroaryl having from 5 to 14 ring members, which contains 1 to 5 heteroatoms as ring members, wherein heteroatoms are selected from N, O and S, and in which heteroaryl is unsubstituted or optionally substituted once or more, independently of each other, by R², in which R² is

hydrogen,

-(C₁-C₄)-alkyl,

-OH,

-O-(C₁-C₄)-alkyl,

halogen, or

Application Ser. No.: 10/613,588

Filing Date: July 3, 2003

Examiner: Huang, Evelyn Mei

~~-N(R³)-R⁴, in which R³ and R⁴ are, independently of each other, hydrogen atom or -(C₁-C₄)-alkyl, then at least one of the following applies:~~

B is not a covalent bond or -(C₁-C₄)-alkylene;

D is not heteroaryl, heterocycle, ~~-(C₆-C₁₄)-aryl, -(C₃-C₆)-cycloalkyl, or~~

~~X and Z are not -(C₁-C₄)-alkyl, OH, O-(C₁-C₄)-alkyl, or halogen.~~

12. (original) The composition as claimed in claim 11, wherein

A is -(C₁-C₃)-alkyl, in which alkyl is straight-chain or branched and is optionally substituted, once or more, independently of each other, by

-O-R¹, or

-C(O)-OR¹, in which R¹ is

hydrogen,

-(C₁-C₃)-alkyl, or

-CF₃

fluoroalkyl of the formula -C_nH_xF_y or fluoroalkoxy of the formula -OC_nH_xF_y, wherein n is an integer from 1 to 3, x is an integer from 0 to 6, y is an integer from 1 to 7 and sum of x and y is 2n + 1,

B is a covalent bond or O,

D is phenyl or naphthyl, in which phenyl or naphthyl is unsubstituted or substituted, once or more, independently of each other, by R², in which R² is

fluorine, chlorine or bromine,

-OH,

-CF₃,

-SR¹, in which R¹ is defined as above,

Application Ser. No.: 10/613,588

Filing Date: July 3, 2003

Examiner: Huang, Evelyn Mei

-(C₁-C₄)-alkyl
-O-(C₁-C₂)-alkyl or
-N(R³)-R⁴, in which R³ and R⁴ are, independently of each other, hydrogen atom or -(C₁-C₃)-alkyl,

heteroaryl selected from the group consisting of pyridyl, furanyl, pyrrolyl, isoxazolyl, benzofuranyl, benzothiophenyl, quinolinyl, isoquinolinyl, quinoxalinyl and thiophenyl, in which heteroaryl is unsubstituted or substituted, once or more, independently of each other, by R², in which R² is defined as above or

-(C₄-C₆)-cycloalkyl, in which cycloalkyl is unsubstituted or substituted, once or more, independently of each other, by R², and R² is defined as above, or

B-D is ((CH₂)_a-Y-R³), in which a is an integer from 1 to 2, Y is O and R³ is -(C₁-C₃)-alkyl, and

R is hydrogen,

-(C₁-C₃)-alkyl, or
-phenyl-(C₁-C₃)-alkyl, and

X and Z are identical or different and are, independently of each other, hydrogen, -C(O)-O(C₁-C₃)alkyl, -OCH₃, -N(CH₃)₂ or halogen.

13. (original) The composition as claimed in claim 11, wherein the compound of the formula I is selected from the group consisting of:

5-pyridin-2-yl-3-trifluoromethyl-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(2-trifluoromethyl-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(3-trifluoromethyl-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(4-trifluoromethyl-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,

Application Ser. No.: 10/613,588

Filing Date: July 3, 2003

Examiner: Huang, Evelyn Mei


1,3-dimethyl-5-(3-trifluoromethylphenyl)-1H-pyrazolo[4,3-c]-isoquinoline,
5-phenyl-3-trifluoromethyl-1H-pyrazolo[4,3-c]isoquinoline,
1,3-dimethyl-5-(3-trifluoromethylphenyl)-1H-pyrazolo[4,3-c]-isoquinoline,
1,3-dimethyl-5-(2,6-difluorophenyl)-1H-pyrazolo[4,3-c]-isoquinoline,
1-benzyl-5-cyclohexyl-3-methyl-1H-pyrazolo[4,3-c]-isoquinoline,
1-benzyl-5-naphthyl-3-methyl-1H-pyrazolo[4,3-c]-isoquinoline,
5-methoxymethyl-3-methyl-1H-pyrazolo[4,3-c]-isoquinoline,
7-methoxycarbonyl-3-methyl-5-phenyl-1H-pyrazolo[4,3-c]-isoquinoline,
7-methoxycarbonyl-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]-
isoquinoline,
7-dimethylamino-3-methyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,
7-dimethylamino-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]-isoquinoline,
6-dimethylamino-3-methyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,
6-dimethylamino-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]-isoquinoline,
8-dimethylamino-3-methyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,
8-dimethylamino-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]-isoquinoline,
1,3-dimethyl-5-(3-methyl-thiophen-2-yl)-1H-pyrazolo[4,3-c]-isoquinoline,
3-methyl-5-phenyl-9-trifluoromethyl-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-pyridin-2-yl-9-trifluoromethyl-1H-pyrazolo[4,3-c]-isoquinoline,
and
3-methyl-5-(2,3,4,5,6-pentafluoro-phenyl)-1H-pyrazolo[4,3-c]-isoquinoline.

14. (currently amended) A method of treating a disease condition associated with inflammation comprising administering to a patient suffering from said disease condition a therapeutically effective amount of a compound of formula (I):

Application Ser. No.: 10/613,588

Filing Date: July 3, 2003

Examiner: Huang, Evelyn Mei



(I)

or a stereoisomeric form or a pharmaceutically acceptable salt of the compound of the formula I, wherein

A is $-(C_1-C_6)$ -alkyl, in which alkyl is straight-chain or branched and is optionally substituted, once or more, independently of each other, by

- O-R¹ or
- C(O)-OR¹,
- C(O)-NR¹R¹,
- C(O)-NR¹-SO₂R¹,
- NR¹R¹,
- CN, in which R¹ is
 - hydrogen,
 - $-(C_1-C_6)$ -alkyl,
 - $-(C_8-C_{14})$ aryl or

fluoroalkyl of the formula $-C_nH_xF_y$ or fluoroalkoxy of the formula $-OC_nH_xF_y$, wherein n is an integer from 1 to 6, x is an integer from 0 to 12, y is an integer from 1 to 13 and sum of x and y is $2n + 1$,

- O-R¹,
- SR¹,
- S(O)-R¹
- S(O)₂-R¹

Application Ser. No.: 10/613,588

Filing Date: July 3, 2003

Examiner: Huang, Evelyn Mei

-C(O)-OR¹,

fluoroalkyl of the formula -C_nH_xF_y or fluoroalkoxy of the formula -OC_nH_xF_y, wherein n is an integer from 1 to 6, x is an integer from 0 to 12, y is an integer from 1 to 13 and sum of x and y is 2n + 1,

-C(O)-NR¹R¹,

-C(O)-NR¹-SO₂R¹,

-NR¹R¹,

-CN,

monocyclic or polycyclic heteroaryl having from 5 to 14 ring members, which contains 1 to 5 heteroatoms as ring members, wherein heteroatoms are selected from N, O and S, and in which heteroaryl is unsubstituted or optionally substituted once or more, independently of each other, by

R², in which R² is

-(C₁-C₄)-alkyl,

-OH,

-O-(C₁-C₄)-alkyl,

halogen,

-N(R³)-R⁴, in which R³ and R⁴ are, independently of each other, hydrogen atom or -(C₁-C₄)-alkyl,

fluoroalkyl of the formula -C_nH_xF_y or fluoroalkoxy of the formula -OC_nH_xF_y, wherein n is an integer from 1 to 4, x is an integer from 0 to 8, y is an integer from 1 to 9 and sum of x and y is 2n + 1,

-CN,

-SR¹,

-S(O)-R¹,

-S(O)₂-R¹ or

Application Ser. No.: 10/613,588

Filing Date: July 3, 2003

Examiner: Huang, Evelyn Mei

$-\text{C}(\text{O})-\text{NR}^1\text{R}^1$,

$-(\text{C}_3-\text{C}_6)$ -cycloalkyl, in which cycloalkyl is unsubstituted or substituted, once or more, independently of each other, by R^2 , and R^2 is defined as above,

monocyclic or bicyclic heterocycle having from 5 to 12 ring members, which contains 1 to 5 heteroatoms as ring members, wherein heteroatoms are selected from N, O and S, and in which heterocycle is unsubstituted or substituted, once or more, independently of each other, by R^2 , and R^2 is defined as above,

B is a covalent bond,

$-\text{C}=\text{CR}^1-$,

$-\text{C}\equiv\text{C}-$,

$-\text{O}(\text{CH}_2)_a-$, in which a is an integer from 1 to 4,

O, S, NR^2 , $-\text{C}(\text{O})-$, $-\text{NR}^2-\text{C}(\text{O})-$, $-\text{C}(\text{O})-\text{NR}^2-$, $-\text{NR}^2-\text{SO}_2-$, $-\text{SO}_2-\text{NR}^2-$, $-\text{NR}^2-\text{C}(\text{O})-\text{NR}^2-$, and R^2 is defined as above, or

$-(\text{C}_1-\text{C}_4)$ -alkylene, in which alkylene is straight-chain or branched and is optionally substituted, once or more, independently of each other, by R^1 , and R^1 is defined as above,

D is $-(\text{C}_1-\text{C}_6)$ -alkyl, in which alkyl is straight-chain or branched and is optionally substituted, once or more, independently of each other, by R^1 , and R^1 is defined as above,

monocyclic or polycyclic heteroaryl having from 5 to 14 ring members, which contains 1 to 5 heteroatoms as ring members, wherein heteroatoms are selected from N, O and S, and in which

Application Ser. No.: 10/613,588

Filing Date: July 3, 2003

Examiner: Huang, Evelyn Mei

heteroaryl is unsubstituted or is substituted once or more,
independently of each other, by R^2 , and R^2 is defined as above,

monocyclic or bicyclic heterocycle having from 5 to 12 ring
members, which contains 1 to 5 heteroatoms as ring members,
wherein heteroatoms are selected from N, O and S, and in which
heterocycle is unsubstituted or substituted, once or more,
independently of each other, by R^2 , and R^2 is defined as above,

-(C_6 - C_{14})-aryl, in which aryl is unsubstituted or substituted,
once or more, independently of each other, by R^2 , and R^2 is defined
as above, or

-(C_3 - C_8)-cycloalkyl, in which cycloalkyl is unsubstituted or
substituted, once or more, independently of each other, by R^2 , and R^2
is defined as above, or

B-D is hydrogen,
halogen,

fluoroalkyl of the formula $-C_nH_xF_y$ or fluoroalkoxy of the
formula $-OC_nH_xF_y$, wherein n is an integer from 1 to 4, x is an integer
from 0 to 8, y is an integer from 1 to 9 and sum of x and y is $2n + 1$,

-(CH_2)_a-Y- R^3 , in which a is an integer from 1 to 4, Y is O, S, NR^2 , and
 R^3 is

-(C_1 - C_6)-alkyl,
-(C_6 - C_{14})-aryl,
-(C_3 - C_8)-cycloalkyl, and

R is hydrogen,

Application Ser. No.: 10/613,588

Filing Date: July 3, 2003

Examiner: Huang, Evelyn Mei

-(C₁-C₆)-alkyl, or
-(C₆-C₁₄)-aryl-(C₁-C₈)-alkyl, in which aryl is unsubstituted or substituted, once or more, independently of each other, by R², and R² is defined as above, and

X and Z are identical or different and are, independently of each other selected from:

hydrogen atom,
-(C₁-C₄)-alkyl,
-OH,
-O-(C₁-C₄-alkyl),
halogen,
fluoroalkyl of the formula -C_nH_xF_y or fluoroalkoxy of the formula -OC_nH_xF_y, wherein n is an integer from 1 to 6, x is an integer from 0 to 12, y is an integer from 1 to 13 and sum of x and y is 2n + 1,
-C(O)-OR¹,
-C(O)-NR¹R¹,
-C(O)-NR¹-SO₂R¹,
-NR¹R¹,
-NR¹-C(O)-NR¹R¹,
-NR¹-C(O)-R¹,
-NR¹-C(O)-OR¹,
-O-C(O)-NR¹R¹,
-CN,
-SR¹,
-S(O)-R¹,
-S(O)₂-R¹,
-S(O)₂-NR¹R¹,
-NR¹-SO₂-R¹, in which R¹ is as defined above,

Application Ser. No.: 10/613,588

Filing Date: July 3, 2003

Examiner: Huang, Evelyn Mei

heterocycle having from 5 to 12 ring members, in which heterocycle is unsubstituted or substituted, once or more, independently of each other, by R^2 , and R^2 is defined as above, or

-(C₃-C₆)-cycloalkyl, in which cycloalkyl is unsubstituted or substituted, once or more, independently of each other, by R^2 , and R^2 is defined as above,

with the proviso that when A is -(C₁-C₆)-alkyl, in which alkyl is straight-chain or branched and is optionally substituted, once or more, independently of each other, by

-O-R¹ or
-C(O)-OR¹, in which R¹ is
hydrogen,
-(C₁-C₆)-alkyl,

-O-R¹, in which R¹ is as defined in this proviso,

-C(O)-OR¹, in which R¹ is as defined in this proviso, or

monocyclic or polycyclic heteroaryl having from 5 to 14 ring members, which contains 1 to 5 heteroatoms as ring members, wherein heteroatoms are selected from N, O and S, and in which heteroaryl is unsubstituted or optionally substituted once or more, independently of each other, by R^2 , in which R^2 is

hydrogen,
-(C₁-C₄)-alkyl,
-OH,
-O-(C₁-C₄)-alkyl,
halogen, or

-N(R³)-R⁴, in which R³ and R⁴ are, independently of each other, hydrogen atom or -(C₁-C₄)-alkyl, then

B is not a covalent bond or -(C₁-C₄)-alkylene.

15. (original) The method as claimed in claim 14 wherein

A is -(C₁-C₃)-alkyl, in which alkyl is straight-chain or branched and is optionally substituted, once or more, independently of each other, by

Application Ser. No.: 10/613,588

Filing Date: July 3, 2003

Examiner: Huang, Evelyn Mei

-O-R¹, or
-C(O)-OR¹, in which R¹ is
hydrogen,
-(C₁-C₃)-alkyl, or
-CF₃

fluoroalkyl of the formula -C_nH_xF_y or fluoroalkoxy of the
formula -OC_nH_xF_y, wherein n is an integer from 1 to 3, x is an
integer from 0 to 6, y is an integer from 1 to 7 and sum of x and y is
2n + 1,

B is a covalent bond or O,

D is phenyl or naphthyl, in which phenyl or naphthyl is
unsubstituted or substituted, once or more, independently of each
other, by R², in which R² is

fluorine, chlorine or bromine,
-OH,
-CF₃,
-SR¹, in which R¹ is defined as above,
-(C₁-C₄)-alkyl
-O-(C₁-C₂)-alkyl or
-N(R³)-R⁴, in which R³ and R⁴ are, independently of
each other, hydrogen atom or -(C₁-C₃)-alkyl,

heteroaryl selected from the group consisting of
pyridyl, furanyl, pyrrolyl, isoxazolyl, benzofuranyl, benzothiophenyl,
quinoliny, isoquinoliny, quinoxaliny and thiophenyl, in which
heteroaryl is unsubstituted or substituted, once or more,
Independently of each other, by R², in which R² is defined as above or

Application Ser. No.: 10/613,588

Filing Date: July 3, 2003

Examiner: Huang, Evelyn Mei

-(C₄-C₆)-cycloalkyl, in which cycloalkyl is unsubstituted or substituted, once or more, independently of each other, by R², and R² is defined as above, or

B-D is ((CH₂)_a-Y-R³, in which a is an integer from 1 to 2, Y is O and R³ is -(C₁-C₃)-alkyl, and

R is hydrogen,

-(C₁-C₃)-alkyl, or

-phenyl-(C₁-C₃)-alkyl, and

X and Z are identical or different and are, independently of each other, hydrogen, -C(O)-O(C₁-C₃)alkyl, -OCH₃, -N(CH₃)₂ or halogen.

16. (original) The method as claimed in claim 14, wherein the compound of formula (I) is selected from the group consisting of:

5-pyridin-2-yl-3-trifluoromethyl-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(2-trifluoromethyl-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(3-trifluoromethyl-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(4-trifluoromethyl-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
1,3-dimethyl-5-(3-trifluoromethylphenyl)-1H-pyrazolo[4,3-c]-isoquinoline,
5-phenyl-3-trifluoromethyl-1H-pyrazolo[4,3-c]isoquinoline,
1,3-dimethyl-5-(3-trifluoromethylphenyl)-1H-pyrazolo[4,3-c]-isoquinoline,
1,3-dimethyl-5-(2,6-difluorophenyl)-1H-pyrazolo[4,3-c]-isoquinoline,
1-benzyl-5-cyclohexyl-3-methyl-1H-pyrazolo[4,3-c]-isoquinoline,
1-benzyl-5-naphthyl-3-methyl-1H-pyrazolo[4,3-c]-isoquinoline,
5-methoxymethyl-3-methyl-1H-pyrazolo[4,3-c]-isoquinoline,
7-methoxycarbonyl-3-methyl-5-phenyl-1H-pyrazolo[4,3-c]-isoquinoline,
7-methoxycarbonyl-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]-isoquinoline,

Application Ser. No.: 10/613,588

Filing Date: July 3, 2003

Examiner: Huang, Evelyn Mei

7-dimethylamino-3-methyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,
7-dimethylamino-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]-isoquinoline,
6-dimethylamino-3-methyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,
6-dimethylamino-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]-isoquinoline,
8-dimethylamino-3-methyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,
8-dimethylamino-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]-isoquinoline,
1,3-dimethyl-5-(3-methyl-thiophen-2-yl)-1H-pyrazolo[4,3-c]-isoquinoline,
3-methyl-5-phenyl-9-trifluoromethyl-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-pyridin-2-yl-9-trifluoromethyl-1H-pyrazolo[4,3-c]-isoquinoline,
and
3-methyl-5-(2,3,4,5,6-pentafluoro-phenyl)-1H-pyrazolo[4,3-c]-isoquinoline.

17. (original) The method as claimed in claim 14, wherein the compound of formula (I) is selected from the group consisting of:

5-benzo[b]thiophene-2-yl-3-methyl-1H-pyrazolo[4,3-c]isoquinoline,
7-bromo-3-methyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,
7-bromo-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]isoquinoline,
5-(3-chloro-benzo[b]thiophen-2-yl)-3-methyl-1H-pyrazolo[4,3-c]-
isoquinoline,
5-(2-chloro-pyridin-3-yl)-3-methyl-1H-pyrazolo[4,3-c]isoquinoline,
5-(6-chloro-pyridin-3-yl)-3-methyl-1H-pyrazolo[4,3-c]isoquinoline,
3-ethyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,
3-ethyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]isoquinoline,
3-ethyl-5-pyridin-3-yl-1H-pyrazolo[4,3-c]isoquinoline,
5-furan-2-yl-3-methyl-1H-pyrazolo[4,3-c]isoquinoline,
7-methoxy-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(3-methyl-benzofuran-2-yl)-1H-pyrazolo[4,3-c]-isoquinoline,
3-methyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,
6-chloro-3-methyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,
6-fluoro-3-methyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,

Application Ser. No.: 10/613,588

Filing Date: July 3, 2003

Examiner: Huang, Evelyn Mei

8-chloro-3-methyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,
8-fluoro-3-methyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,
9-chloro-3-methyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,
9-fluoro-3-methyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,
3,9-dimethyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(2-methyl-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(3-methyl-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(4-methyl-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(2-bromo-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(3-bromo-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(2-chloro-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(4-chloro-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(2,4-dichloro-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(3,4-dichloro-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(2,6-dichloro-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(2-fluoro-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(4-fluoro-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(2,4-difluoro-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(2,6-difluoro-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
6-chloro-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]isoquinoline,
6-fluoro-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]isoquinoline,
8-chloro-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]isoquinoline,
8-fluoro-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]isoquinoline,
9-chloro-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]isoquinoline,
9-fluoro-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]isoquinoline,
3,9-dimethyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(1-methyl-1H-pyrrol-2-yl)-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-quinolin-2-yl-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-quinolin-3-yl-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-quinoxalin-2-yl-1H-pyrazolo[4,3-c]isoquinoline,

Application Ser. No.: 10/613,588
Filing Date: July 3, 2003
Examiner: Huang, Evelyn Mei

3-methyl-5-thiophen-2-yl-1H-pyrazolo[4,3-c]isoquinoline, and
3-methyl-5-(3-methyl-thiophen-2-yl)-1H-pyrazolo[4,3-c]isoquinoline.

18. (original) The method as claimed in claim 14, wherein the disease condition is selected from the group consisting of multiple sclerosis, atherosclerosis, inflammatory bowel disease, Alzheimer's disease, stroke and diabetes.
19. (original) The method as claimed in claim 18, wherein the disease condition is multiple sclerosis.
20. (original) The method as claimed in claim 18, wherein the disease condition is atherosclerosis.
21. (original) The method as claimed in claim 18, wherein the disease condition is inflammatory bowel disease.
22. (original) The method as claimed in claim 18, wherein the disease condition is Alzheimer's disease.
23. (original) The method as claimed in claim 18, wherein the disease condition is stroke.
24. (original) The method as claimed in claim 18, wherein the disease condition is diabetes.